



# Olivatis® 20

## Applicative guidelines

### INTRODUCTION

**Olivatis® 20** is a liquid, O/W emulsifier designed for cold process applications. It generates stable emulsions but requires a correct applicative procedure. The aim of these guidelines is to help formulators to use **Olivatis® 20** successfully.

**Olivatis® 20** is compatible with all kinds of oil-soluble ingredients such as esters, triglycerides and silicones.

### GENERAL EMULSIFICATION PROCEDURE

#### 1. USE WITH HIGH-SHEAR MIXER

If a high-shear mixer is available, **Olivatis® 20** can be mixed with oils and the achieved phase can be added to the water phase under high shear.

#### 2. USE WITHOUT HIGH-SHEAR MIXER

**Olivatis® 20** must be pre-dispersed in water before the addition of fats and the required amount of water is usually 5 times the amount of **Olivatis® 20** (e.g. a formulation containing 2,5% of **Olivatis® 20** should be dispersed in 15% of water). The dispersion can be slightly rheopectic (viscosity increases increasing the speed of stirrer) therefore speed must be reduced in case of thickening.

After the dispersion of **Olivatis® 20**, add fats and oils under stirring and make a primary emulsion.

Once the emulsion is homogenous, add the rest of water containing stabilizers, preservatives and other water-soluble ingredients.

#### 3. MANUFACTURING OF GEL CREAM WITH CARBOMED 1030

**OLIVATIS® 20** is indicated for the formulation of gel cream with **Carbomed 1030**. To obtain a perfect gel cream, **Carbomed 1030** must be added at the end of the formulation, as the last ingredient and the quantity of **Carbomed 1030** required can vary from 0,25 to 0,5% depending on the final thickness required.

**Carbomed 1030** does not produce lumps so it is possible to add it without worries directly to the emulsion formed with the methods described previously. The addition of **Carbomed 1030** must be done by keeping the emulsion under stirring and adding the neutralizer directly to the emulsion. This last operation must be done after the powder has been completely incorporated into the emulsion and considering that 1,33 g of NaOH 30% or 1,5 g of TEA are required for each gram of **Carbomed 1030**. After the addition of neutralizer emulsion must be stirred slowly until the gel is homogeneous.



## REQUIRED CONCENTRATION

Usually, better results are achieved with a ratio of 1 part of **Olivatis® 20** with 2 parts of oil phase. The ratio may change depending on the nature of the oil used in the formulation.

## STABILIZERS

Fluid emulsions can be stabilized with ordinary thickening agents such as xanthan gum, hydroxypropyl guar, sclerotium gum, etc. Inorganic thickeners such as magnesium aluminium silicate, are compatible and useful as stabilizers too. It is recommended to disperse gums into the water phase.

The addition of slight amounts (0,5-1%) of sorbitan monooleate increase the emulsion stability. Long- chain fatty alcohols, such as octyldodecanol, also increase the emulsion stability.

## INCOMPATIBILITIES

Large amounts of electrolytes (>1% of NaCl) can interfere with the emulsification while large amounts of glycols (> 5% propylene glycol, butylene glycol, etc.) can affect emulsion stability.

The quaternary ammonium compounds are generally compatible. However, given the large number of available compounds and the difference in the chemical structure, compatibility must be assessed case by case.



## FORMULATION EXAMPLE WITH HIGH-SHEAR MIXER

### BODY MILK BASE FORMULATION

INGREDIENTS	%	PHASE
Olivatis® 20	2,5	I
Octyl Stearate	3,0	I
Dicaprylyl Ether	3,0	I
Tocopheryl Acetate	0,5	I
Phenoxyethanol, Ethylhexylglycerin	1,0	II
Glycerin	1,0	II
Xanthan Gum	0,2	II
Water	q.s. to 100,0	II

### PROCEDURE

- Phase I**  
 Mix all ingredients in a separate vessel.
- Phase II**  
 Disperse Xanthan Gum in the rest of water, add the preservative (Phenoxyethanol, Ethylhexylglycerin) and Glycerin, stir until all ingredients are dissolved.
- Under high-shear mixer add phase I to phase II, homogenize for at least 30 minutes.
- Results**  
 White fluid milk  
 pH 7,0



## FORMULATION EXAMPLE WITHOUT HIGH-SHEAR MIXER

### BODY MILK BASE FORMULATION

INGREDIENTS	%	PHASE
Olivatis® 20	2,5	I
Octyl Stearate	3,0	I
Dicaprylyl Ether	3,0	I
Tocopheryl Acetate	0,5	I
Phenoxyethanol, Ethylhexylglycerin	1,0	II
Glycerin	1,0	II
Xanthan Gum	0,2	II
Water for primary emulsion	15,0	I
Water	q.s. to 100,0	II

### PROCEDURE

- Phase I**  
 Disperse **Olivatis® 20** in water for primary emulsion, stir until it is completely dispersed.  
 Separately, mix Octyl Stearate, Dicaprylyl Ether, Tocopheryl Acetate and add to the **Olivatis® 20** emulsion, keep stirring.
- Phase II**  
 Disperse Xanthan Gum in the rest of water, add the preservative (Phenoxyethanol, Ethylhexylglycerin) and Glycerin, stir until all ingredients are dissolved.
- Mix phase I and phase II under stirring, check pH.
- Results**  
 White fluid milk  
 pH 7,0

Revision 01 Date: 18.12.2023

This information is for illustration purposes only and based upon our knowledge and experience at the time of writing. We do not grant any warranty of any kind, express or implied, including merchantability or fitness for a particular use, with respect to such information or its application. Use of this information is at users' discretion and risk. Nothing herein relieves users from carrying out their own suitability determinations and tests. Users are responsible for observing all local laws and regulations relevant to their products, uses or processes. We do not assume any liability in connection with their products and its use. The preceding information and suggestions are also provided without warranty of non-infringement as to the intellectual property rights of third parties, and shall not be construed as any inducement to infringe the rights of third parties.